

# HOW TO TAKE OXBRYTA® (voxelotor) 500 MG TABLETS



FOR PATIENTS 12 YEARS AND UP, AS WELL AS PATIENTS 4 TO LESS THAN 12 WHO WEIGH 88 LBS (40 KG) OR MORE\*

ARELYS ... Mother, avid traveler, and actual Oxbryta patient

Oxbryta comes in 2 different dosage forms. Your healthcare provider will decide which dosage form you should take. **This dosing card is for use with Oxbryta 500 mg Tablets only**, which are swallowed whole. **Do not** cut, crush, or chew the tablets.

## STEPS FOR TAKING YOUR MEDICINE

- 1 Visually inspect the pills to ensure it is the correct medication. The Oxbryta 500 mg Tablets are film-coated, light yellow, oval shaped, and marked with "GBT 500" on one side. (Please see photo below.)
- 2 Swallow tablets whole once a day or exactly as your healthcare provider tells you.
  - **Do not** cut, crush, or chew the tablets

RECOMMENDED DOSE	NUMBER OF OXBRYTA 500 MG TABLETS
1,000 mg	2
1,500 mg	3
2,000 mg	4
2,500 mg	5

- 3 Take with or without food.



## IMPORTANT INFORMATION

- Take Oxbryta exactly as your healthcare provider tells you
- **Do not** change your dose or stop taking Oxbryta unless your healthcare provider tells you to
- Take the full prescribed dose of Oxbryta 1 time each day
- Your healthcare provider may change your dose if needed
- Your healthcare provider may also prescribe a medicine called hydroxyurea during treatment with Oxbryta
- **If you miss a dose or do not take the entire dose of Oxbryta**, skip that dose and return to your normal dosing schedule the next day
- Do not take St. John's wort during treatment with Oxbryta

### HOW TO STORE OXBRYTA

- Store Oxbryta at 68°F to 86°F (20°C to 30°C)
  - Oxbryta comes in a child-resistant package
  - The bottle contains a desiccant to help keep your medicine dry (protect it from moisture) and polyester coil. Do not eat
- Keep Oxbryta and all medicines out of the reach of children.**

\*Based on their ability to swallow tablets.

For full information on how you should take Oxbryta, please see the [Patient Prescribing Information on Oxbryta.com](#). If you have any questions, please talk to your healthcare provider.

## INDICATION

### What is OXBRYTA?

OXBRYTA is a prescription medicine used for the treatment of sickle cell disease in adults and children 4 years of age and older.

It is not known if OXBRYTA is safe and effective in children with sickle cell disease below 4 years of age.

This indication is approved under accelerated approval based on increase in hemoglobin (Hb). Continued approval for this indication may be contingent upon verification and description of clinical benefit in confirmatory trial(s).

## IMPORTANT SAFETY INFORMATION

**Do not take OXBRYTA** if you or your child have had an allergic reaction to voxelotor or any of the ingredients in OXBRYTA. See the end of the patient leaflet for a list of the ingredients in OXBRYTA.

**Please see additional Important Safety Information on page 2 and full [Prescribing Information on Oxbryta.com](#).**



## INDICATION

### What is OXBRYTA?

OXBRYTA is a prescription medicine used for the treatment of sickle cell disease in adults and children 4 years of age and older.

It is not known if OXBRYTA is safe and effective in children with sickle cell disease below 4 years of age.

This indication is approved under accelerated approval based on increase in hemoglobin (Hb). Continued approval for this indication may be contingent upon verification and description of clinical benefit in confirmatory trial(s).

## IMPORTANT SAFETY INFORMATION

**Do not take OXBRYTA** if you or your child have had an allergic reaction to voxelotor or any of the ingredients in OXBRYTA. See the end of the patient leaflet for a list of the ingredients in OXBRYTA.

**Before taking OXBRYTA, tell your healthcare provider about all of your medical conditions, including if you or your child:**

- have liver problems
- are pregnant or plan to become pregnant. It is not known if OXBRYTA can harm your unborn baby.
- are breastfeeding or plan to breastfeed. It is not known if OXBRYTA can pass into your breastmilk and if it can harm your baby. Do not breastfeed during treatment with OXBRYTA and for at least 2 weeks after the last dose.

**Tell your healthcare provider about all the medicines you or your child take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Some medicines may affect how OXBRYTA works. OXBRYTA may also affect how other medicines work and may affect the results of certain blood tests. Keep a list of all your medicines and show it to your healthcare provider.

### What should I avoid while taking OXBRYTA?

Do not take St. John's wort during treatment with OXBRYTA.

### What are the possible side effects of OXBRYTA?

**OXBRYTA can cause serious side effects, including:**

**Serious allergic reactions.** Tell your healthcare provider or get emergency medical help right away if you get:

- rash
- shortness of breath (difficult breathing)
- hives
- swelling of the face

**The most common side effects of OXBRYTA include:**

- headache
- nausea
- diarrhea
- rash or hives
- stomach-area (abdominal) pain
- fever

**The most common side effects of OXBRYTA in children ages 4 to less than 12 years of age include:**

- fever
- stomach-area (abdominal) pain
- vomiting
- diarrhea
- rash
- headache

These are not all the possible side effects of OXBRYTA.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. You may also report side effects to Global Blood Therapeutics at 1-833-428-4968 (1-833-GBT-4YOU).

**Keep OXBRYTA and all medicines out of the reach of children.**

**Please see full Prescribing Information on Oxbryta.com.**

