

HOW TO TAKE OXBRYTA® (voxelotor) 300 MG TABLETS FOR ORAL SUSPENSION

FOR PATIENTS 4 YEARS AND OLDER BASED ON
WEIGHT AND ABILITY TO SWALLOW TABLETS




HAJAR ... Student, artist, and actual Oxbryta patient

Oxbryta comes in 2 different dosage forms. Your healthcare provider will decide which dosage form you should take. **This dosing card is for use with Oxbryta 300 mg Tablets for Oral Suspension only, which should first be mixed in room temperature clear drink such as drinking water or clear soda before taking. Do not** swallow whole, cut, crush, or chew the tablets for oral suspension.

Visually inspect the pills to ensure it is the correct medication. (Please see photo below.)

HOW TO PREPARE

- A teaspoon
- A small cup
- A clear drink (For example: drinking water or clear soda)
- Oxbryta 300 mg Tablets for Oral Suspension, which are light yellow, round shaped, and marked with "300 D" on one side



Approximate
size



0.43 in. wide

IMPORTANT INFORMATION

- Take Oxbryta exactly as your healthcare provider tells you
- **Do not** change your dose or stop taking Oxbryta unless your healthcare provider tells you to
- Take the full prescribed dose of Oxbryta 1 time each day
- Your healthcare provider may change your dose if needed
- Your healthcare provider may also prescribe a medicine called hydroxyurea during treatment with Oxbryta
- Take Oxbryta with or without food
- **If you miss a dose or do not take the entire dose of Oxbryta**, skip that dose and return to your normal dosing schedule the next day
- Do not take St. John's wort during treatment with Oxbryta

Please see full Prescribing Information and Instructions for Use for complete information on Dosage and Administration.

IMPORTANT SAFETY INFORMATION

Do not take OXBRYTA if you or your child have had an allergic reaction to voxelotor or any of the ingredients in OXBRYTA. See the end of the patient leaflet for a list of the ingredients in OXBRYTA.

Please see additional Important Safety Information on page 4 and full Prescribing Information on Oxbryta.com.

INDICATION

What is OXBRYTA?

OXBRYTA is a prescription medicine used for the treatment of sickle cell disease in adults and children 4 years of age and older.

It is not known if OXBRYTA is safe and effective in children with sickle cell disease below 4 years of age.

This indication is approved under accelerated approval based on increase in hemoglobin (Hb). Continued approval for this indication may be contingent upon verification and description of clinical benefit in confirmatory trial(s).



STEPS FOR TAKING YOUR MEDICINE

1 POUR CLEAR DRINK



Visually inspect the pills to ensure it is the correct medication. Please see photo on prior page.

Be sure your hands are clean and dry, then pour room temperature clear drink in the cup. Use the table below as a guide for the minimum amount of liquid needed for the prescribed dose. You may add more clear drink if needed to mix the tablets.



NUMBER OF 300 MG TABLETS	AMOUNT OF CLEAR DRINK
1	1 tsp (5 mL)
2	2 tsp (10 mL)
3	3 tsp (15 mL)
4	4 tsp (20 mL)
5	5 tsp (25 mL)
7	7 tsp (35 mL)
8	8 tsp (40 mL)

2 ADD TABLET(S)



Add the prescribed number of Oxbryta 300 mg Tablets for Oral Suspension to the cup.

- **Do not** swallow whole, cut, crush, or chew the tablets for oral suspension


3 SWIRL & WAIT



Swirl the cup until the tablet(s) break down in the drink. Take care not to spill. **The tablet(s) will not completely dissolve. You will still see small tablet clumps in the mixture.**

- If you spill any medicine, clean up the spill. Throw away the rest of the prepared medicine and start again

Wait for 1 to 5 minutes before taking or giving the dose.



For full information on how you should take Oxbryta, please see the [Patient Prescribing Information](#) and [Instructions for Use](#) on [Oxbryta.com](#). If you have any questions, please talk to your healthcare provider.

Please see Important Safety Information on page 4 and full [Prescribing Information](#) on [Oxbryta.com](#).



4 SWIRL & DRINK



Swirl the cup again. Take care not to spill. Drink immediately.

- If giving to a child, make sure that the child is upright. Give all the prepared medicine to the child immediately



5 ADD MORE LIQUID & REPEAT



Add 1 or 2 more teaspoons of clear drink to the cup. This is to mix the leftover medicine and to make sure the full dose is taken. Swirl the cup until all the leftover medicine is mixed, and drink immediately. Repeat this step until all the medicine is taken.

- After taking all the medicine, you may drink more water or any type of drink
- Be sure to wash the spoon and cup with warm water and soap



For more details on how to take Oxbryta 300 mg Tablets for Oral Suspension, please see the [Patient Prescribing Information](#) and [Instructions for Use on Oxbryta.com](#).

HOW TO STORE OXBRYTA

- Store Oxbryta at 68°F to 86°F (20°C to 30°C)
- Oxbryta comes in a child-resistant package
- The bottle contains a polyester coil. Do not eat

Keep Oxbryta and all medicines out of the reach of children.

INDICATION

What is OXBRYTA?

OXBRYTA is a prescription medicine used for the treatment of sickle cell disease in adults and children 4 years of age and older.

It is not known if OXBRYTA is safe and effective in children with sickle cell disease below 4 years of age.

This indication is approved under accelerated approval based on increase in hemoglobin (Hb). Continued approval for this indication may be contingent upon verification and description of clinical benefit in confirmatory trial(s).

IMPORTANT SAFETY INFORMATION

Do not take OXBRYTA if you or your child have had an allergic reaction to voxelotor or any of the ingredients in OXBRYTA. See the end of the patient leaflet for a list of the ingredients in OXBRYTA.

Before taking OXBRYTA, tell your healthcare provider about all of your medical conditions, including if you or your child:

- have liver problems
- are pregnant or plan to become pregnant. It is not known if OXBRYTA can harm your unborn baby.
- are breastfeeding or plan to breastfeed. It is not known if OXBRYTA can pass into your breastmilk and if it can harm your baby. Do not breastfeed during treatment with OXBRYTA and for at least 2 weeks after the last dose.

Tell your healthcare provider about all the medicines you or your child take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Some medicines may affect how OXBRYTA works. OXBRYTA may also affect how other medicines work and may affect the results of certain blood tests. Keep a list of all your medicines and show it to your healthcare provider.

What should I avoid while taking OXBRYTA?

Do not take St. John's wort during treatment with OXBRYTA.

What are the possible side effects of OXBRYTA?

OXBRYTA can cause serious side effects, including:

Serious allergic reactions. Tell your healthcare provider or get emergency medical help right away if you get:

- rash
- shortness of breath (difficult breathing)
- hives
- swelling of the face

The most common side effects of OXBRYTA include:

- headache
- nausea
- diarrhea
- rash or hives
- stomach-area (abdominal) pain
- fever

The most common side effects of OXBRYTA in children ages 4 to less than 12 years of age include:

- fever
- stomach-area (abdominal) pain
- vomiting
- diarrhea
- rash
- headache

These are not all the possible side effects of OXBRYTA.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. You may also report side effects to Global Blood Therapeutics at 1-833-428-4968 (1-833-GBT-4YOU).

Keep OXBRYTA and all medicines out of the reach of children.

Please see full Prescribing Information on [Oxbryta.com](https://www.oxbryta.com).

